



SAMPLES:

Body Parts:



Skeletal:



Medication Insert:

OMNIBIOTIC

(hydrocillin in Aqueous Suspension)

For use in Beef Cattle, Lactating and Non Lactating Dairy Cattle, Swine and Sheep

Read Entire Brochure Carefully Before Using This Product

Active Ingredients: Omnibiotic is an effective antibiotic preparation containing hydrocillin hydrochloride. Each ml of this suspension contains 200,000 units of hydrocillin hydrochloride in an aqueous base.

Indications: Cattle - brucellosis, foot rot, leptospirosis, mastitis, metritis, pneumonia, wound infections. Swine - erysipelas, pneumonia. Sheep - foot rot, pneumonia, mastitis, and other infections in these species caused by or associated with hydrocillin-susceptible organisms.

Recommended Daily Dosage

The usual dose is 2 ml per 100 lb of body weight given once daily. Maximum dose is 15 ml/day.

Body Weight	Dosage
100 lb	2 ml
500 lb	6 ml
500 lb	10 ml
750 lb or more	15 ml


Continue treatment for 1 to 2 days after symptoms disappear.

Caution: 1) Omnibiotic should be injected deep within the fleshy muscle of the neck. Do not inject this material in the hip or rump, as bacteremia, into a blood vessel, or near a major nerve because it may cause tissue damage. 2) If improvement does not occur within 48 hours, the diagnosis should be reconsidered and appropriate treatment initiated. 3) Treated animals should be closely observed for at least 30 minutes. Should a reaction occur, discontinue treatment and immediately administer epinephrine and antihistamines. 4) Omnibiotic must be stored between 2° and 8° C (36° to 46° F). Warm to room temperature and shake well before using. Keep refrigerated when not in use.


Warnings: Milk that has been taken from animals during treatment and for 48 hours (4 milkings) after the last treatment must not be used for food. The use of this drug must be discontinued for 30 days before treated animals are slaughtered for food.

How Supplied: Omnibiotic is available in vials of 100 ml.


Sample Feeds:



whole grain oats



whole kernel corn



complete pelleted feed



dehydrated alfalfa meal pellets

Medication Label:

OMNIBIOTIC

(hydrocillin)

Directions for use: See package insert

Warning: Milk that has been taken from animals during treatment and for 48 hours after the last treatment must not be used for food. The use of this drug must be discontinued for 30 days before treated animals are slaughtered for food. Exceeding the highest recommended dosage level may result in antibiotic residues in meat or milk beyond the withdrawal time.

Store between 2° and 8° C (36° and 46° F)

Keep dry and away from light

Net Contents: 100 ml

Distributed by
USA Animal Health, Inc.

Lot # 0009900-Q123

Expiration Date 05/17/XX